

PHH harmonic submersions are stable

Monica Alice Aprodu

University Dunărea de Jos, Domneasca 47, 6200, Galati, Romania, Email: Monica.APRODU@ugal.ro

Abstract. We prove that PHH harmonic submersions are (weakly) stable.

A harmonic map between Riemann manifolds is called (*weakly*) *stable* if the Hessian of the energy functional is (semi) positive definite, see, for example [7], Chapter 5. In particular, an energy-minimizing map is stable. Lichnerowich has proved in 1970 (see [4]) that holomorphic maps between Kähler manifolds are (weakly) stable; away from these particular mappings, we do not dispose of many other examples of harmonic maps which are (weakly) stable.

In the joint paper [1], we have introduced a class of harmonic maps, defined on a Riemann manifold, with value in a Kähler manifold, called *PHH harmonic maps* which have a behaviour somewhat similar to that of holomorphic maps. Holomorphic maps between Kähler manifolds are typical examples of PHH harmonic maps, but examples of different flavour have been found in [2].

The aim of this paper is to prove that PHH harmonic submersions are actually (weakly) stable, yet another property which relates maps in this class to holomorphic maps (compare to [3]).

We start by recalling some basic notions and facts from [1], and [5], see also [3]. We start with a map $\varphi : (M^m, g_M) \rightarrow (N^{2n}, J_N, g_N)$, defined on a Riemann manifold with value in a Kähler manifold. For any point $x \in M$, we denote $d\varphi_x^* : T_{\varphi(x)}N \rightarrow T_xM$ the adjoint of the tangent map $d\varphi_x : T_xM \rightarrow T_{\varphi(x)}N$. The map φ is called PHWC at x (*pseudo-horizontally weakly conformal*) if and only if $d\varphi_x \circ d\varphi_x^*$ commutes with $J_{N, \varphi(x)}$. Naturally, φ is called PHWC if it is PHWC at any point $x \in M$. If φ is PHWC at x , we say that φ is PHH at x (*pseudo-horizontally homothetic*) if and only if

$$d\varphi_x ((\nabla_v^M d\varphi_x^*(J_N Y))_x) = J_{N, \varphi(x)} d\varphi_x ((\nabla_v^M d\varphi_x^*(Y))_x)$$

for any horizontal tangent vector $v \in T_xM$, and any vector field Y , locally defined in a neighbourhood of $\varphi(x)$, where ∇^M is the Levi-Civita connection on M (the Levi-Civita connections on N is denoted by ∇^N , and the induced connection in the bundle $\varphi^{-1}TN$ is denoted by $\tilde{\nabla}$). The map φ is called PHH if it is PHH at any point x of M . It is easy to see that a PHWC map is PHH if and only if

$$d\varphi(\nabla_X^M d\varphi^*(J_N Y)) = J_N d\varphi(\nabla_X^M d\varphi^*(Y)),$$

for any horizontal vector field X on M and any vector field Y on N .

The two conditions PHWC and PHH can be seen in terms of the almost complex structure on the horizontal bundle, defined by $J_H = d\varphi^{-1} \circ J_N \circ d\varphi$. J_H is the restriction of an f -structure on M , also denoted by J_H , which vanishes on the vertical distribution. The PHWC condition is the compatibility condition of J_H with the metric on M , while pseudo-horizontally homothetic condition is equivalent to J_H being parallel in horizontal directions.

Suppose next that the map φ is harmonic and submersive, and M is compact. In this case, we know from Theorem 2.1 (a), Proposition 3.1 and Proposition 3.3 in [1] that the fibres of φ are minimal submanifolds. Recall next that the stability of harmonic maps is controlled by a condition on the Hessian of the energy-functional:

$$H(E)_\varphi(V, V) \geq 0,$$

for all V a section of the bundle $\varphi^{-1}TN$. The Hessian is computed by (see [7], pp. 155)

$$H(E)_\varphi(V, W) = \int_M g_N(J_\varphi V, W) v_M,$$

for all V, W sections in $\varphi^{-1}TN$, where v_M is the volume form on M , and J_φ is a second order selfadjoint elliptic differential operator acting on sections of $\varphi^{-1}TN$ in the following way. Denote by NR the curvature tensor field on N , and let $\{\varepsilon_1, \dots, \varepsilon_m\}$ be an orthogonal vector frame on M , and V be a section in $\varphi^{-1}TN$. Then

$$J_\varphi V := - \sum_{i=1}^m \frac{1}{\|\varepsilon_i\|^2} \left(\tilde{\nabla}_{\varepsilon_i} \tilde{\nabla}_{\varepsilon_i} - \tilde{\nabla}_{\nabla_{\varepsilon_i} \varepsilon_i} \right) V - \sum_{i=1}^m \frac{1}{\|\varepsilon_i\|^2} {}^NR(V, d\varphi(\varepsilon_i)) d\varphi(\varepsilon_i).$$

The second-order elliptic differential operator

$$\bar{\Delta}_\varphi := - \sum_{i=1}^m \frac{1}{\|\varepsilon_i\|^2} \left(\tilde{\nabla}_{\varepsilon_i} \tilde{\nabla}_{\varepsilon_i} - \tilde{\nabla}_{\nabla_{\varepsilon_i} \varepsilon_i} \right)$$

is called the *rough Laplacian* of φ , cf. [7], pp. 155. The second sum which appears in the formula defining the *Jacobi operator* J_φ is denoted by $\mathcal{R}_\varphi$, so $J_\varphi = \bar{\Delta}_\varphi - \mathcal{R}_\varphi$. One of the useful properties of the rough Laplacian, which will be constantly used in the sequel is the following, cf. [7], pp. 156.

Proposition 1 *The rough Laplacian $\bar{\Delta}_\varphi$ satisfies*

$$\int_M g_N(\bar{\Delta}_\varphi V, W) v_M = \int_M g_N(\tilde{\nabla} V, \tilde{\nabla} W) v_M = \int_M g_N(V, \bar{\Delta}_\varphi W) v_M,$$

where V and W are sections on $\varphi^{-1}TN$, and

$$g_N(\tilde{\nabla} V, \tilde{\nabla} W) = \sum_{i=1}^m \frac{1}{\|\varepsilon_i\|^2} g_N(\tilde{\nabla}_{\varepsilon_i} V, \tilde{\nabla}_{\varepsilon_i} W).$$

After these preparations, we arrive at the statement of the main result of this paper.

Theorem 2 *Let (M^m, g_M) be a compact Riemann manifold, (N^{2n}, J_N, g_N) be a Kähler manifold, and $\varphi : M \rightarrow N$ be a harmonic PHH submersion. Then φ is (weakly) stable.*

Proof. As in the proof of Theorem 4.1 of [1], we choose a (local) frame $\{e_1, \dots, e_n, J_N e_1, \dots, J_N e_n\}$ in $\varphi^{-1}TN$ such that the system $\{d\varphi^*(e_1), \dots, d\varphi^*(e_n), d\varphi^*(J_N e_1), \dots, d\varphi^*(J_N e_n)\}$ is an orthogonal frame in the horizontal distribution. We also choose $\{u_1, \dots, u_s\}$ an orthonormal basis for the vertical distribution. We denote $E_i = d\varphi^*(e_i)$, and $E'_i = d\varphi^*(J_N e_i)$, for all $i = 1, \dots, n$.

With this notation, we apply the same strategy of proof as in [7], pp. 172, Theorem 3.2.

For V a section in $\varphi^{-1}TN$, we apply Proposition 1, and compute:

$$H(E)_\varphi(V, V) = \int_M g_N(\tilde{\nabla} V, \tilde{\nabla} V) v_M - \int_M g_N(\mathcal{R}_\varphi V, V) v_M.$$

By definition

$$\begin{aligned} g_N(\tilde{\nabla} V, \tilde{\nabla} V) &= \sum_{i=1}^n \left(\frac{1}{\|E_i\|^2} g_N(\tilde{\nabla}_{E_i} V, \tilde{\nabla}_{E_i} V) + \frac{1}{\|E'_i\|^2} g_N(\tilde{\nabla}_{E'_i} V, \tilde{\nabla}_{E'_i} V) \right) \\ &\quad + \sum_{j=1}^s g_N(\tilde{\nabla}_{u_j} V, \tilde{\nabla}_{u_j} V). \end{aligned}$$

Analogous to the operator used in the proof of Theorem 3.2, Chapter 5, [7], we define, for any $V \in \Gamma(\varphi^{-1}TN)$, the operator $DV \in \Gamma(\varphi^{-1}TN \otimes \mathcal{H}^*)$, where $\mathcal{H}$ is the horizontal distribution on M , by

$$DV(X) := \tilde{\nabla}_{J_H X} V - J_N \tilde{\nabla}_X V,$$

for any X a horizontal vector field on M .

Next, we compute

$$\begin{aligned} g_N(DV, DV) &= \sum_{i=1}^n \left\{ \frac{1}{\|E_i\|^2} g_N(DV(E_i), DV(E_i)) \right. \\ &\quad \left. + \frac{1}{\|E'_i\|^2} g_N(DV(E'_i), DV(E'_i)) \right\}. \end{aligned}$$

Since $J_H E_i = E'_i$, $J_H E'_i = -E_i$, and $\|E_i\| = \|E'_i\|$, we obtain

$$\begin{aligned} g_N(DV, DV) &= 2 \sum_{i=1}^n \frac{1}{\|E_i\|^2} \left(g_N(\tilde{\nabla}_{E_i} V, \tilde{\nabla}_{E_i} V) + g_N(\tilde{\nabla}_{E'_i} V, \tilde{\nabla}_{E'_i} V) \right. \\ &\quad \left. - 2g_N(\tilde{\nabla}_{E'_i} V, J_N \tilde{\nabla}_{E_i} V) \right) \end{aligned}$$

Therefore

$$\begin{aligned} \int_M \left(g_N(J_\varphi V, V) - \frac{1}{2} g_N(DV, DV) \right) v_M &= \sum_{i=1}^n \int_M \frac{1}{\|E_i\|^2} \left(2g_N(\tilde{\nabla}_{E'_i} V, J_N \tilde{\nabla}_{E_i} V) \right. \\ &\quad \left. - g_N(^N R(V, d\varphi(E_i)) d\varphi(E_i), V) \right. \\ &\quad \left. - g_N(^N R(V, d\varphi(E'_i)) d\varphi(E'_i), V) \right) v_M. \end{aligned}$$

Next, taking into account the identities $d\varphi(E'_i) = J_N d\varphi(E_i)$, $d\varphi(E_i) = -J_N d\varphi(E'_i)$, and the basic properties of the curvature tensor field $^N R$, we obtain

$$^N R(V, d\varphi(E_i)) d\varphi(E_i) + ^N R(V, d\varphi(E'_i)) d\varphi(E'_i) = J_N ^N R(d\varphi(E_i), d\varphi(E'_i)) V,$$

and thus

$$\begin{aligned} &\int_M \left(g_N(J_\varphi V, V) - \frac{1}{2} g_N(DV, DV) \right) v_M \\ &= \sum_{i=1}^n \int_M \frac{1}{\|E_i\|^2} \left(2g_N(\tilde{\nabla}_{E'_i} V, J_N \tilde{\nabla}_{E_i} V) - g_N(J_N ^N R(d\varphi(E_i), d\varphi(E'_i)) V, V) \right) v_M. \end{aligned}$$

We compute

$$\begin{aligned} -g_N(J_N ^N R(d\varphi(E_i), d\varphi(E'_i)) V, V) &= g_N(^N R(d\varphi(E_i), d\varphi(E'_i)) V, J_N V) \\ &= g_N(\tilde{\nabla}_{E_i} \tilde{\nabla}_{E'_i} V - \tilde{\nabla}_{E'_i} \tilde{\nabla}_{E_i} V - \tilde{\nabla}_{[E_i, E'_i]} V, J_N V) \\ &= E_i g_N(\tilde{\nabla}_{E'_i} V, J_N V) - E'_i g_N(\tilde{\nabla}_{E_i} V, J_N V) \\ &\quad - g_N(\tilde{\nabla}_{\nabla_{E_i} E'_i} V, J_N V) + g_N(\tilde{\nabla}_{\nabla_{E'_i} E_i} V, J_N V) \\ &\quad - g_N(\tilde{\nabla}_{E'_i} V, \tilde{\nabla}_{E_i} J_N V) + g_N(\tilde{\nabla}_{E_i} V, \tilde{\nabla}_{E'_i} J_N V). \end{aligned}$$

Similarly to [7], pp. 180, we define a $\mathcal{C}^\infty$ function ϕ on M by the formula:

$$\begin{aligned} \phi &:= \sum_{i=1}^n \frac{1}{\|E_i\|^2} \left(E_i g_N(\tilde{\nabla}_{E'_i} V, J_N V) - E'_i g_N(\tilde{\nabla}_{E_i} V, J_N V) \right. \\ &\quad \left. - g_N(\tilde{\nabla}_{\nabla_{E_i} E'_i} V, J_N V) + g_N(\tilde{\nabla}_{\nabla_{E'_i} E_i} V, J_N V) \right). \end{aligned}$$

Since

$$g_N(\tilde{\nabla}_{E_i} V, \tilde{\nabla}_{E'_i} J_N V) = -g_N(\tilde{\nabla}_{E_i} J_N V, \tilde{\nabla}_{E'_i} V),$$

we have

$$\int_M \left(g_N(J_\varphi V, V) - \frac{1}{2} g_N(DV, DV) \right) v_M = \int_M \phi v_M.$$

The proof of the Theorem will be concluded if we prove

$$\int_M \phi v_M = 0.$$

For this, we use Green's formula. We choose X a horizontal vector field on M defined by the property:

$$g_M(X, Y) = g_N(\tilde{\nabla}_{J_H Y} V, J_N V),$$

for any vector field Y on M , and we prove $\operatorname{div}(X) = \phi$. Indeed, since the fibres of φ are minimal, and X is horizontal, it follows:

$$\operatorname{div}(X) = \sum_{i=1}^n \frac{1}{\|E_i\|^2} \left(g_M(E_i, \nabla_{E_i} X) + g_M(E'_i, \nabla_{E'_i} X) \right).$$

Next,

$$\begin{aligned} \operatorname{div}(X) &= \sum_{i=1}^n \frac{1}{\|E_i\|^2} \left(E_i g_M(E_i, X) - g_M(\nabla_{E_i} E_i, X) + E'_i g_M(E'_i, X) - g_M(\nabla_{E'_i} E'_i, X) \right) \\ &= \sum_{i=1}^n \frac{1}{\|E_i\|^2} \left(E_i g_N(\tilde{\nabla}_{J_H E_i} V, J_N V) - g_N(\tilde{\nabla}_{J_H \nabla_{E_i} E_i} V, J_N V) \right) \\ &\quad + \sum_{i=1}^n \frac{1}{\|E_i\|^2} \left(E'_i g_N(\tilde{\nabla}_{J_H E'_i} V, J_N V) - g_N(\tilde{\nabla}_{J_H \nabla_{E'_i} E'_i} V, J_N V) \right). \end{aligned}$$

By the PHH condition, we have $J_H \nabla_{E'_i} E'_i = -(\nabla_{E'_i} E_i)^h$, and $J_H \nabla_{E_i} E_i = (\nabla_{E_i} E'_i)^h$, where by $(\cdot)^h$ we denoted the horizontal component of $(\cdot)$, so,

$$\operatorname{div}(X) = \phi.$$

We proved

$$\int_M g_N(J_\varphi V, V) v_M = \frac{1}{2} g_N(DV, DV) v_M \geq 0.$$

Remark 3 *Our result improves the main result of [6], provided that the source manifold is compact (condition which is not needed in [6]).*

Acknowledgements. This work was financed by a NATO fellowship. The author is grateful to the Fourier Institute in Grenoble for hospitality, and to J. C. Wood for some useful remarks on an early version of the manuscript.

References

- [1] M. A. Aprodu, M. Aprodu, V. Brînzănescu, A class of harmonic maps and minimal submanifolds. *Int. J. Math.* **11** (2000), 1177-1191.
- [2] M. A. Aprodu, M. Aprodu, Implicitly defined harmonic PHH submersions. *Manuscripta Math.* **100** (1999), 103-121.
- [3] D. Burns, F. Burstall, P. De Bartolomeis, J. Rawnsley, Stability of harmonic maps of Kähler manifolds. *J. Differential Geom.* **30** (1989), 579-594.
- [4] A. Lichnerowicz, Applications harmoniques et variétés kählériennes. *Symp. Math.* III (Bologna 1970), 341-402.
- [5] E. Loubeau, Pseudo Harmonic Morphisms. *Int. J. Math.* **7** (1997), 943-957.
- [6] S. Montaldo, Stability of harmonic morphisms to a surface, *Int. J. Math.* **9** (1998), 865-875.
- [7] K. Urakawa, Calculus of variations and harmonic maps. *Transl. Math. Monographs* Vol. 132. AMS Providence, Rhode Island: 1993.